

F.G.6





FIG.7

FIG.8


1.6x10<sup>6</sup>/12well N=5

Culture time: 3,7,14,21day
Asc-2P : 0, 0.1, 1, 5mM



Culture time (day)

Hydroxyproline(pg)







FIG. 10








FIG. 12




FIG.13




FIG. 14





FIG. 15









FIG.17





















FIG.20





Cultured for 7 days (Hunziker EB. JBJS 1996) remove superficial zone chondroitinase ABC digested with FIG.23






FIG. 22








FIG.25



**FIG.26** 



FIG.27










FIG.29





membrana synovialis derived artificial tissue













FIG.32





HE staining

FIG.34









38/46









FIG.39


Ratio of initial diameter







**FIG.43** 



**FIG.44** 







FIG.45







contact projection A

$$H=\frac{F}{A}=\frac{F}{k_l h_p^2}$$

$$E = \left[\frac{dF}{dh}\right]_{F_{max}} \frac{1 - \nu^2}{2 \cdot k_2 \cdot h_{pmax}}$$

$$h_p = h_r + 0.25(h_{\text{max}} - h_r)$$

F: load

A: contact projection area

hp: contact depthk1k2: shape conflictFmax: Maximum load

hmax: Maximum displacement hr: point at which tangential

line intersects

dF/dh: Gradient of tangential line of load removal curve

ν: Poisson's ratio